Bays, Estuaries, and Riparian Habitats

♦ *Facilitators*:

Rainer Hoenicke, San Francisco Estuary Institute

♦ *Rapporteurs*:

Susan Pufahl, Monterey Bay National Marine Sanctuary

♦ *Participants*:

Ross Clark, California Coastal Commission

Ted Grosholz, University of California at Davis

Fred LaCaro, State Water Resources Control Board

Dave Paradies, Bay Foundation of Morro Bay

Holly Price, Monterey Bay National Marine Sanctuary

Don Potts, University of California at Santa Cruz

Victoria Seidman, California State Parks

Mark Stephenson, Calif. Dept. of Fish and Game / Moss Landing Marine Labs

Kerstin Wasson, Elkhorn Slough Foundation and

Elkhorn Slough National Estuarine Research Reserve

Fred Watson, California State University Monterey Bay

Karen Worcester, California Regional Water Quality Control Board

Susanne Worcester, California State University Monterey Bay

♦ *Priority Issues*:

- Habitat Modification
- Pollution
- Biodiversity / Ecosystem Integrity
- Invasive Species
- Global, Large Scale Human Impacts
- ♦ *Priority Questions* (for each issue above):
 - What is the extent of impact?
 - Which resources are affected?
 - How can we best monitor effectiveness of restoration, prevention and preservation efforts?
 - What are the opportunities for restoration, prevention and preservation efforts?
 - What are historic conditions and carrying capacities?
 - Is status changing?
 - What is seasonal, annual, long-term and spatial variability?
- ♦ Characterization of Priority Issues or Questions:

Habitat Modification - Habitat Loss

Parameters - Area of marsh, wetland and riparian habitats

Methods - Aerial surveys/GIS analysis for habitat acreage, field surveys to ground truth above and measure habitat structure (with biodiversity)

Spatial Scale - From satellite imagery to quarter meter quadrats with emphasis on developed coastal areas

Temporal Scale - Every two to five years, indefinitely

Frequency - Winter and summer seasons

Existing Data/Programs - USGS hydrologic quad resources (NHD), numerous habitat maps

Habitat Modification - Bottlenecks, Sinks, Rivermouths, Confluences and Harbors

Parameters - Pollutant loading and biological impacts, sources, sediment and flow dynamics *Methods* - Chemical, tissue, and sediment samples; plume characterizations with aerial photos; species composition at all levels

Spatial Scale - Specific regions or habitats of concern

Temporal Scale - Five years rotations and after storm events (depends on parameter), indefinitely *Frequency* - Parameter dependent

Existing Data/Programs - NPDES, Mussel Watch, Municipalities

Is the number of invading species changing in the Sanctuary?

Parameters - Species richness of exotics, abundance of a few know pest species

Methods - Will vary by habitat and taxon

Spatial Scale - Entire MBNMS

Temporal Scale - Indefinitely

Frequency - Twice per year

Existing Data/Programs - Wasson et al. summary; Grosholtz detailed studies

Additional Comments - Must look beyond just estuaries and identify the source; a key is to detect invasions early for best chance of eradication

What are the impacts of invading species on specific communities and habitats?

Parameters - Species richness, abundance, distribution, and behavior of target species, selected physical or chemical parameters

Methods - Will vary by habitat and taxon

Spatial Scale - Entire MBNMS

Temporal Scale - Indefinitely

Frequency - Twice per year

Existing Data/Programs - Grosholtz impact of green crabs

Additional Comments - Must look at links between invasive and native species

What are estuarine ecosystem components?

Parameters - Communities structure, biodiversity, patchiness, and productivity in salt marshes, eelgrass, and mudflats

Methods - Use the diversity of bird species as indicators; sediment samples, species lists, invertebrates and microbial communities, density of dominant plants; remote sensing, GIS and local counts; comprehensive baseline surveys, inventory and mapping

Spatial Scale - Entire Sanctuary, down to individual habitats

Temporal Scale - Indefinitely

Frequency - Two to four times per year

Additional Comments - Long-term community involvement for funding and help in comprehensive sampling

Pollutants

Parameters - Nutrients, pathogens, sediments, organophospates, chlorinated hydrocarbons; bioaccumulation, toxicity analysis and basic physical parameters

Methods - Standard methods and new technologies (e.g., for nitrates)

Spatial Scale - Rivermouths and watersheds to help identify sources, at risk nearshore habitats Temporal Scale - Indefinitely

Frequency - Monthly and after storm events

Existing Data/Programs – Numerous (e.g., MBNMS WQPP, CDFG, CRWQCB)

Additional Comments - Initial sampling is needed to identify watershed loading and source allocations contributing to loading